



ISSN: 0975-766X

CODEN: IJPTFI  
Research Article

Available Online through  
[www.ijptonline.com](http://www.ijptonline.com)

**DEVELOPMENT AND VALIDATION OF UV-SPECTROPHOTOMETRIC  
METHOD FOR DETERMINATION OF QUETIAPINE FUMARATE IN  
PURE AND PHARMACEUTICAL FORMULATION**

**Susamma Cicy Eapen, V G Prasanth\*, Sheeja Velayudhan Kutty, T S Jyothi**  
Department of Pharmaceutical Analysis, Grace College of Pharmacy, Kodunthirapully P O,  
Palakkad, India, Pin 678004.

Email: [prasanthvelekkat@gmail.com](mailto:prasanthvelekkat@gmail.com)

Received on 01-10-2011

Accepted on 23-10-2011

**Abstract**

A simple, sensitive, selective, economical and reproducible UV spectrophotometric method has been developed for the quantitative determination of QTF in bulk drug and in pharmaceutical dosage forms. The methods are based on measurement of absorbance of QTF solution in ethanol at 207 nm. Beer's law is obeyed over the linear range 1-5  $\mu\text{g/mL}$  of QTF for the method with apparent molar absorptivity value of  $1434.41281 \text{ L mol}^{-1}\text{cm}^{-1}$ . Limits of quantification (LOQ) and detection (LOD) are also reported. The methods were validated in accordance with the current ICH guidelines. The precision results, expressed by intra-day and inter-day relative standard deviation values, are satisfactory i.e % RSD 100.22% and 99.83 % respectively. The accuracy is also satisfactory (%RSD 0.39) and percentage recoveries are in the range 99.34- 100.11% with the standard deviation of 0.39. Method have excellent linearity and range ( $r^2 = 0.998$ ).

**Keywords:** Quetiapine Fumarate, UV spectrophotometry, molar absorptivity

An atypical antipsychotic, Quetiapine fumarate (2-[2-(4-dibenzo[*b,f*] [1,4]thiazepin-11-yl-1-piperazinyloxy) ethanol fumarate (2:1 salt)) which has a unique receptor-binding profile belonging to a new chemical class, the dibenzothiazepine derivatives<sup>[1]</sup>. Quetiapine is an antagonist at a broad range of neurotransmitter receptors. Quetiapine is used in the treatment of schizophrenia or manic episodes associated with bipolar disorder. As a consequence, there is an increasing demand for new analytical methods for determination of same drug in most economical way. Several HPLC methods for

the determination of QUET have been reported, most of these require ultraviolet detection<sup>[2-4]</sup> as QUET is not electro active, some stability indicating<sup>[5]</sup>, impurity characterizing<sup>[6]</sup>. A HPTLC method has been developed<sup>[7]</sup>. However none of these methods is sensitive enough for determination of the expected drug levels and some of them are time consuming and require complex sample pretreatment or long run times. Some gas chromatography–mass spectrometry (GC–MS) methods have also been employed, however here QUET needs to be derivatized before analysis<sup>[8]</sup>. Some HPLC–MS-MS methods has been published for determination of QUET<sup>[9]</sup>. The goal of our work was to develop an UV spectrophotometric method for determination of QUET in solid dosage form and to use the results for analysis of drug in pharmaceuticals in most economic way, as rapid and effective ways for determination of drugs in sample matrix by spectroscopy is desirable as such no analytical paper is available for the quality control of pharmaceutical formulations containing QUET by spectroscopy.


## Materials & methods

A Shimadzu Model PHARAMASPEC-1800 UV–Vis spectrophotometer with 1.0 cm matched cells was used for the electronic spectral measurements. Quetiapine Fumarate and all other chemicals used were analytical reagent grade (AR grade). Ethanol was used as a solvent in all experimental purpose. QTF pure drug (certified to be 99.85% pure) was kindly provided by Dr. Reddy's Laboratories Ltd Hyderabad, India, as a gift and used as received. Qutipin-100 (100 mg QTF) were manufactured by Sun Pharmaceuticals Ltd, India and purchased .

### I. Solutions

An accurately weighed quantity of 10 mg QUET was transferred into 100 mL volumetric flask with sufficient quantity of ethanol and sonicated. The volume was made up to the mark with ethanol. Aliquots of this standard stock solution (SSS) of QUET was diluted with ethanol and scanned over the UV range of 200-400 nm. A spectrum of drug was drawn out and selected the wavelength 207 nm for the analysis at which drug showed maximum absorbance (**Fig. No.1**).

Figure no 1. Determination of  $\lambda_{max}$  of Quet.



## II. Procedure

### 1. For calibration curve: (study of Beer- Lambert’s Law)

Five mixed standards of QUE having concentrations 1, 2, 3, 4 and 5  $\mu\text{g/mL}$  were prepared from SSS . Individual and overlay absorption spectra were processed at 207 nm (Fig. No 2 and 3). Absorbance at five different standards plotted against concentration and calibration graph forms .(Table no 2).

Figure no.2: Plot of Beers- Lambert’s Law for Quet AT 207 nm.





Figure no. 3: Overlay Spectra of Quet.



## 2. For absorptivity study:

From the SSS, a solution of 5 µg/ml concentration was prepared. Absorbance of such five of QUET standard solution measured and results of absorptivity study out by A1% 1cm. (**Table No.1**)

## 3. Estimation of QUET in tablet formulation sample:

Twenty tablets were weighed accurately and powdered. Powder equivalent to 10 mg<sup>[10]</sup> (Label claim- 100 mg ) was taken and transferred to 100 ml volumetric flask and dissolved in ethanol , sonicate for 10 min. , filtered and further diluted to get a final concentration of 5 µg/mL of QUET (label claim basis). The absorbance of solution is measured at the selected wavelength i.e . 207 nm .( **Table No. 2**)

## III. Result and discussion

The method was accurate , simple , rapid , reliable , sensitive , reproducible and economical . The wavelength 207 nm was selected which showed good linearity between concentrations .

## IV Validation of analytical data

The study of Beer's -Lambert's law was checked by preparing standard solutions at 5 different concentrations and the linearity of the calibration graphs and conformity of the UV measurements of the proposed methods to Beer's law were proven by the values of the correlation coefficient (r) of the absorptivity study . The linear range of concentrations for the analysis of QUET was found to be 1-5 µg mL<sup>-1</sup> for UV spectrophotometric method . (**Figure No.2-A & Figure No .2-B**)

The utility of this method was verified by analysis of a recovery assay in the marketed tablet sample . Tablet sample (Label claim- 100mg ) QUET was prepared and processed according to the proposed method . Recoveries were determined by standard addition method (SAM). The mean percentage recoveries of QUET by UV method were found to be 99.68 % . Results represent accuracy by study of recovery. (**Table No.3**)

The reproducibility of this developed method established by study of precision for QUET was determined by five replicates analysis on the tablet sample<sup>[11]</sup>. The mean relative standard deviations were found to be 100.27 % for UV spectrophotometric method. (**Table No.2**)

Correlation coefficient for UV spectrophotometric method was found to be 0.998 of QUET was found to be linear. (**Figure No.2**)

The Ruggedness of the proposed method checked by means of two parameters i.e. Intraday & Interday<sup>[12]</sup>. For intra –day inter –day study, standard deviation and relative standard deviations were found to be 0.68724, 0.685732 % and 0.72437, 0.7255953 % respectively. (**Table No.4**)

#### IV. Conclusion

Validation parameters complies, the applied spectrophotometric methods of analysis are simple, sensitive, accurate and satisfactorily capable for determination of QUET in tablet formulation with reproducible specific results. The linear concentration range of preordain elaborated method were observed wider (**Table no-5**). In addition, the analyses by proposed method is cheaper and economic too. Thus, proposed UV spectrophotometric method is applicable for the quality control and routine analysis and may also be proposed for determination from biological fluids or other solid dosage form containing same drugs.

**Table No.1: Absorptivity (1%, 1cm) Values of Quet At 207 nm.**

Sl. No.	Conc. g/100mL	Abs.	A (1%, 1cm)*
1	0.0005072	0.720	1419.55836
2	0.0005683	0.734	1444.02912
3	0.0005003	0.712	1423.14611
4	0.0005126	0.743	1449.47327
5	0.0005098	0.732	1435.85721
<b>Mean</b>			<b>1434.41281</b>
<b>± S.D.</b>			<b>1.06196</b>
<b>% RSD</b>			<b>0.074</b>

**Table No. 2: Estimation of Quet In Tablet Formulation.**

Sl. No.	Wt. Of tablet powder taken mg(label claim 100mg)	Abs. at 207 nm	Amount of drug per tablet (mg)	% Purity
1	21.2	0.720	99.44	99.44
2	21.34	0.734	100.71	100.71
3	21.06	0.712	98.99	98.99
4	21.4	0.743	101.66	101.66
5	21.32	0.732	100.53	100.53
<b>Mean</b>			<b>100.27</b>	<b>100.27</b>
<b>± SD</b>			<b>1.06309</b>	<b>1.06309</b>
<b>% RSD</b>			<b>1.06023</b>	<b>1.06023</b>
<b>Variance</b>			<b>1.13018</b>	<b>1.13018</b>

**Table no 2: Calibration Curve.**

Sl.no	Concentration ( $\mu\text{g/ml}$ )	Absorbance
1	1	0.138
2	2	0.267
3	3	0.428
4	4	0.577
5	5	0.712

**Table no 3: Determination of Accuracy By Percentage Recovery Method.**

Drug	Tablet amount (mg)	Amount of pure drug added (mg)	Level of addition (%)	Total Drug estimate d (mg)	Amount of pure drug recoverd (mg)	% recovery
QTF	21	8	80	18.13	8.09	99.34
	21	10	100	20.22	10.01	100.11
	21	12	120	22.31	11.95	99.58
<b>Mean</b>						<b>99.68</b>
<b><math>\pm</math>SD</b>						<b>0.39402</b>
<b>%RSD</b>						<b>0.39528</b>
<b>Variance</b>						<b>0.15525</b>

**Table No.4: Results of Estimation In Intra-Day Studies And Inter-Day Studies.**

Sl. No	Intra-day			Inter-day		
	Hour	Wt. of tablet powder taken (mg)	% Label Claim	Day	Wt. of tablet powder taken (mg)	% of Labeled Claim
1	0 hr	21.21	99.44	Day-1	21.41	100.65
2	3 hr		100.53	Day-2		99.44
3	6 hr		100.71	Day-3		99.4
<b>Mean</b>			<b>100.22</b>	<b>Mean</b>		<b>99.83</b>
<b><math>\pm</math> SD</b>			<b>0.68724</b>	<b><math>\pm</math> SD</b>		<b>0.72436</b>
<b>% RSD</b>			<b>0.68573</b>	<b>% RSD</b>		<b>0.72559</b>
<b>Variance</b>			<b>0.47230</b>	<b>Variance</b>		<b>0.52469</b>

**Table no-5: Validation Parameters.**

SI No	Parameter	Result
1	Absorption maxima (nm)	207
2	Linearity range ( $\mu\text{g/ml}$ )	1-5
3	Slope	0.143
4	Intercept	0.006
5	Correlation coefficient ( $r^2$ )	0.999
6	Molar absorptivity	1434.41281
7	Accuracy (% recovery)	99.68
8	Precision	100.22% (intra day precision) and 99.83% (inter day precision)
9	LOD ( $\mu\text{g/ml}$ )	0.02
10	LOQ ( $\mu\text{g/ml}$ )	0.12

**References**

1. The Complete Drug Reference (Monograph on CD-ROM), Pharmaceutical Press, London, 2003. 5. P.C. Davis, J. Wong, O. Gefvert, Journal of Pharmaceutical and Biomedical Analysis, 20, 1999, 271–282
3. R. Mandrioli, S. Fanali, A. Ferranti, M.A. Raggi, Journal of Pharmaceutical and Biomedical Analysis, 30, 2002, 969–977.
4. C. Frahnert, M.L. Rao, K. Grasmaeder, Journal of Chromatography, B 794, 2003, 35–47.
5. R.H. Pullen, K.M. Palermo, M.A. Curtis, Journal of Chromatography. , 573, 1992, 49.
6. <http://rasayanjournal.com/View-documentdetails/> 80 A validated stability indicating- HPLC-method for the determination of related substances in quetiapine fumarate. htm
7. Fathalla Belal, Amina Elbrashy, Manal Eid, Jenny Jeehan Nasr, Stability-Indicating HPLC Method for the Determination of Quetiapine: Application to Tablets and Human Plasma, Journal of Liquid Chromatography & Related Technologies, 9 (31), 2008 ,1283 – 1298.
8. Ch. Bharathi, K. J. Prabahar, Ch. S. Prasad, M. Srinivasa Rao, G. N. Trinadhachary, V. K. Handa, R. Dandala\*, A. Naidu, Identification, isolation, synthesis and characterization of impurities of Quetiapine Fumarate, Govi-Verlag , 63, 2008, 1, 14-19.
9. S.R. Dhaneshwar\*, N.G. Patre, and M.V. Mahadik, Stability-Indicating HPTLC Method for Quantitation of Quetiapine Fumarate in the Pharmaceutical Dosage Form, Acta chromatographica, 1(21), 2009, 83–93.
10. D.T. Anderson, K.L. Fritz, Journal of analytical toxicology, 24, 2000, 300–304.

11. Borkar Bhaskar Hiranman\*, Vidhate Sandip, Lohiya R. T., Umekar M. J, Spectrophotometric determination of an atypical antipsychotic compound in pharmaceutical formulation ,International Journal of ChemTech Research CODEN( USA): IJCRGG ISSN : 0974-4290 Vol.1, No.4, pp 1153-1161, Oct-Dec 2009
12. K. Basavaiah\*, N. Rajendraprasad, P.J. Ramesh and K.B. Vinay, Sensitive ultraviolet spectrophotometric determination of quetiapine fumarate in pharmaceuticals, Thai J. Pharm. Sci. 34 (2010) 146-154
13. Bagade S.B.\*, Narkhede S.P., Nikam D. S., Sachde C. K., Spectrophotometric method for determination of Quetiapine fumarate in two different dose tablets International Journal of ChemTech Research CODEN ( USA): IJCRGG ISSN : 0974-4290 Vol.1, No.4, pp 898-904, Oct-Dec 2009.

**Corresponding Author:**

**V G Prasanth**

**Email:** [prasanthvelekkat@gmail.com](mailto:prasanthvelekkat@gmail.com)